

Annual Review of Pharmacology and Toxicology
Age-Related Perioperative
Neurocognitive Disorders:
Experimental Models and
Druggable Targets

Odmara L. Barreto Chang,¹ Katherine L. Possin,²
and Mervyn Maze^{1,3}

¹Department of Anesthesia and Perioperative Care, University of California San Francisco, San Francisco, California, USA; email: Mervyn.maze@ucsf.edu

²Memory and Aging Center, Department of Neurology, and Global Brain Health Institute, University of California San Francisco, San Francisco, California, USA

³Center for Cerebrovascular Research, University of California San Francisco, San Francisco, California, USA

ANNUAL
REVIEWS **CONNECT**

www.annualreviews.org

- Download figures
- Navigate cited references
- Keyword search
- Explore related articles
- Share via email or social media

Annu. Rev. Pharmacol. Toxicol. 2023. 63:321–40

First published as a Review in Advance on
September 13, 2022

The *Annual Review of Pharmacology and Toxicology* is
online at pharmtox.annualreviews.org

<https://doi.org/10.1146/annurev-pharmtox-051921-112525>

Copyright © 2023 by the author(s). This work is licensed under a Creative Commons Attribution 4.0 International License, which permits unrestricted use, distribution, and reproduction in any medium, provided the original author and source are credited. See credit lines of images or other third-party material in this article for license information.



Keywords

perioperative neurocognitive disorders, postoperative delirium, neuroinflammation, cognitive assessment, perioperative brain health, care pathways

Abstract

With the worldwide increase in life span, surgical patients are becoming older and have a greater propensity for postoperative cognitive impairment, either new onset or through deterioration of an existing condition; in both conditions, knowledge of the patient's preoperative cognitive function and postoperative cognitive trajectory is imperative. We describe the clinical utility of a tablet-based technique for rapid assessment of the memory and attentiveness domains required for executive function. The pathogenic mechanisms for perioperative neurocognitive disorders have been investigated in animal models in which excessive and/or prolonged postoperative neuroinflammation has emerged as a likely contender. The cellular and molecular species involved in postoperative neuroinflammation are the putative targets for future therapeutic interventions that are efficacious and do not interfere with the surgical patient's healing process.

INTRODUCTION

As the population ages, an increasing number of older patients will undergo surgical procedures (1). Advanced age is a major risk factor for postoperative cognitive impairment, resulting in prolonged hospital stay, loss of independence, higher health-care costs, and increased morbidity and mortality (2–4). Among older adults, the relative risk of developing delirium increases with each added year of age (5), becoming particularly prevalent in surgical patients over 75 years old (6). Understanding the pathogenesis of this common postoperative complication will help identify those older surgical patients for whom both preventative and restorative interventions may be required. In this review, we describe the clinical condition and best practices for diagnosis and treatment and describe how preclinical models have established putative mechanisms that constitute druggable targets for testing the efficacy and safety of interventions.

PERIOPERATIVE NEUROCOGNITIVE DISORDERS

In recent years, the nomenclature for cognitive impairment in the perioperative period has been revised to align with the terminology used for cognitive disorders listed in the *Diagnostic and Statistical Manual of Mental Health Disorders* (7) (DSM-5) (Figure 1). Perioperative neurocognitive disorder (PND) is the most common complication in older surgical patients, with an estimated 1.6–6.4 million Americans suffering from PND each year (8). PND describes new cognitive impairment or deterioration of existing impairment identified in the period immediately prior to surgery and concluding 12 months after surgery (9) (Figure 1). Cognitive impairment within hours after surgery is often classified as emergence excitation or postanesthesia care unit delirium (9, 10). During the postoperative interval, the most common cognitive impairment is postoperative delirium (POD), which can occur in up to 40% of patients older than 60 years (8). POD occurs up to 1 week postprocedure or until discharge from the hospital (whichever occurs first) (9). POD is characterized by fluctuating changes in attention, mental status, or level of consciousness. Irrespective of whether POD occurred, older surgical patients can manifest delayed neurocognitive recovery if new impairment, or increased impairment that was temporally associated with the surgical period, has not resolved within 30 days. If this impairment is present 31 days to 12 months after surgery, it is defined as postoperative neurocognitive disorder (postoperative NCD), which can occur in up to 15% of older patients (10).

It is important to emphasize that outside of the surgical context, NCDs frequently occur in older adults (11) and are a well-established risk factor for PNDs (12–15). The key difference between mild NCD [also called mild cognitive impairment (MCI)] and major NCD (also called dementia) is that in major NCD, the cognitive decline significantly interferes with activities of




Figure 1

The diagram presents the continuum of the classification of perioperative neurocognitive disorder (PND). Preoperatively, patients are cognitively normal or have a neurocognitive disorder (NCD). After surgery, the cognitive impairment is sometimes classified as emergence excitation or postanesthesia care unit (PACU) delirium while recovering from anesthesia. While patients are in the hospital, cognitive impairment is classified as postoperative delirium (POD). The nomenclature changes to delayed neurocognitive disorder for the period up to 30 days after surgery. After that period and up to 12 months afterward, the cognitive impairment is considered a postoperative NCD. Figure created by Dr. Caroline Prioleau and Sabrina Erloff.

daily living (9, 16). In 2020, the estimated US prevalence of dementia was 6.07 million (11.3% of the population aged 65 or older), and the estimated prevalence of MCI was 12.23 million (11). The most common cause of NCD is Alzheimer's disease (AD), followed by cerebrovascular disease and Lewy body disease (i.e., Parkinson's disease dementia and dementia with Lewy bodies); most NCD patients have brain changes associated with more than one cause of dementia (17). Brief cognitive assessments are useful for detecting cognitive impairment, but a more in-depth evaluation is required for differential diagnosis that identifies underlying etiology. Patients with Lewy body disease are at a particularly high risk for POD (18). While differential diagnosis of NCDs is likely beyond the scope of most preoperative clinical settings, extra precautions should be taken for patients with cognitive impairment and warning signs for Lewy body dementia, including rapid-eye-movement sleep behavior disorder, parkinsonism, visual hallucinations, attention fluctuations, and sensitivity to antipsychotic agents (18).

As extant cognitive impairment is a risk factor for PND (15), professional societies such as the American Society of Anesthesiologists, the American College of Surgeons, and the American Geriatrics Society recommend administration of preoperative cognitive assessments for older patients undergoing surgery (19–21). However, these assessments are rarely performed, resulting in underdetection of preoperative cognitive impairment (22–24) and a lost opportunity for targeted interventions to protect brain health.

PREOPERATIVE COGNITIVE ASSESSMENT

Mild cognitive impairment and dementia are substantially underdiagnosed; diagnosis is delayed until moderate or advanced stages of dementia in 50% or more patients (22–24), with greater delays among racial and ethnic minorities (25). For this reason, preoperative cognitive assessment is critical for detecting cognitive impairment and guiding perioperative care. Several reviews have summarized the brief cognitive assessments typically used to detect cognitive impairment prior to surgery (8, 26). Brief (3–15 min) paper-based assessments such as the Mini Mental State Exam, Montreal Cognitive Assessment, Mini-Cog[®], verbal fluency, and clock drawing test, among others, have been used as preoperative cognitive screening tools. Yet, cognitive assessments continue to be infrequently incorporated into preoperative clinical workflows (27), and a reconsideration of the types of tools that will be successful is needed.

Commonly used preoperative cognitive assessments can be useful to detect dementia, but sensitivity of these tests is lower for MCI (28). Furthermore, most brief cognitive assessments have been inadequately adapted for different language/cultural/educational groups, and the impact of these demographic variables is rarely considered when determining whether a patient is impaired in preoperative settings. As an example, a brief cognitive assessment that was developed and validated for an English-speaking population with moderate to high education might not be appropriate for a recent immigrant who has limited English proficiency and few years of education. Brief cognitive assessments are needed that are accurate for MCI detection across demographic and social groups, and scoring may need adjustment for education and other demographic variables (29, 30).

It has not been well studied which cognitive tests are most accurate for predicting postoperative cognitive disorders, but there is reason to emphasize testing of executive functions and episodic memory. Executive functions refer to a constellation of cognitive abilities that enable and drive adaptive, goal-oriented behavior. They include working memory, complex attention (e.g., inhibition, mental flexibility, and sustained attention), planning, fluency, and self-monitoring (31). Episodic memory refers to the ability to consciously recall personally experienced episodes. The memoranda can be verbal, spatial, object based, or multimodal (32). Executive functions and

episodic memory are typically the first cognitive domains to be affected in preclinical AD (33), and vulnerability in the neural networks underlying these skills may predispose patients to the neuroinflammatory consequences of surgery. In POD, there are diffusion changes that can be detected by magnetic resonance imaging (MRI) in the periventricular, frontal, and temporal white matter (34) and a decrease in frontoparietal connectivity on EEG (35); these regions and structures are crucial for executive functions (36). Furthermore, disconnectivity involving interhemispheric and fronto-thalamo-cerebellar networks, as well as microstructural changes of nuclei involved in limbic and memory functions, predisposes patients to delirium after surgery (37). MRI studies suggest that a reduction of thalamic and hippocampal volumes—crucial regions for episodic memory—may be associated with PND (38).

An example of a brief cognitive assessment that is accurate at detecting MCI and sensitive to the memory and executive function impairment of NCDs is the TabCAT Brain Health Assessment (TabCAT-BHA) (36). Subtests include Favorites (associative memory), Match (executive functions and speed), Animal Fluency (language), and Line Orientation (visuospatial). In addition to subtest scores, a composite score summarizes performance, with an emphasis on the memory and executive tests (39). The subtests rely on brain regions vulnerable to surgical insult and NCDs: Favorites has been associated with medial temporal lobe volumes and tau deposition; Match with frontoparietal and basal ganglia network volumes, frontoparietal tau deposition, and corpus callosum white matter integrity; Line Orientation with right parietal volumes and bilateral occipitoparietal tau deposition; and Animal Fluency with left frontal/temporal atrophy (36, 40–42). Furthermore, TabCAT-BHA has been validated in diverse populations, and regression-based norms can be instantly applied that adjust scores for key sociodemographic factors (39, 43).

The TabCAT-BHA is currently in use to improve the detection of NCDs in primary care (44). If a concern for cognitive impairment has been identified by the primary care provider, a medical assistant administers the 10-min test, and results are integrated into the electronic health record (EHR) to guide the primary care provider on the next steps in evaluation in care. Here, we propose a similar efficient workflow adapted for the surgical setting (**Figure 2**). Prior to surgery, adults ages 65 or older or with risk factors for preexisting cognitive impairment are administered a brief, digital cognitive assessment such as the TabCAT-BHA by clinical staff. Results are integrated into the EHR and flagged if the patient has a high likelihood of cognitive impairment, which is a risk factor for PND.




Figure 2

Proposed care pathway for older adults undergoing surgery. Adults ages 65 or older or with risk factors for preexisting cognitive impairment are administered a brief digital cognitive assessment prior to surgery. Results are integrated into the electronic health record and flagged if the patient has a high likelihood of cognitive impairment. After cognitive impairment has been established, the anesthesiologist or another clinician on the team discloses this finding to the patient with a care partner present. Prehabilitation and delirium care pathways are incorporated for patients at high risk for perioperative neurocognitive disorder (PND). Figure created by Dr. Caroline Prioleau and Sabrina Erlhoff.

MONITORING COGNITIVE TRAJECTORIES

Patients and families commonly complain of postoperative cognitive decline. While accurate monitoring of the cognitive trajectory is important for appropriate diagnosis and treatment, it can be difficult to quantify change and monitor for recovery without a reliable preoperative baseline. Sensitive and reliable tests can establish a neurocognitive baseline to monitor postoperative cognitive dysfunction and trajectories, including decline, recovery, and response to treatment (45–47). Reliable change is measured by defining a change in performance between cognitive scores and adjusting for the expected variance of the test (39). Unfortunately, most paper-based cognitive tests that are used in the preoperative setting are designed for screening but are poor for measuring cognitive change over time. Computer-based cognitive batteries are often better suited for following long-term trajectories because of their better psychometric properties and higher measurement precision (48). The TabCAT-BHA, for example, exhibits excellent long-term stability for patients with MCI and dementia and outperforms standard paper-based tests in this regard (39).

PERIOPERATIVE BRAIN HEALTH CARE

After cognitive impairment has been established, the anesthesiologist or another clinician on the team discloses this finding to the patient with a care partner present (**Figure 2**). The risk of PND is discussed, including confusion, inattention, and memory problems that may occur after the operation. The desirability of proceeding, especially with an elective procedure, should be discussed with the surgical provider and the patient. Next, the care partner should be engaged to participate in medical decision-making, to be present during the postoperative period to assist with orienting and providing a comforting environment for the patient, and to develop a caregiving plan if the patient develops POD.

If time allows prior to surgery, prehabilitation may be considered for older adults at high risk for PND (**Figure 2**). Prehabilitation aims to enhance health, optimize comorbidities, treat sleep disorders, and address modifiable risk factors before surgery via a variety of interventions that typically focus on nutrition, exercise training, and reducing stress and anxiety. Reversible causes of cognitive impairment, including medication side effects, sleep apnea, depression, and other medical conditions, can be treated prior to surgery (49–51). While more research is needed to understand whether prehabilitation reduces the risk of PND (52, 53), there is evidence that prehabilitation programs can decrease hospital length of stay and improve functional capacity (54, 55). Recent research has suggested that cognitive training may offer some benefit (56), although not all studies concur (57). Delirium care pathways should be triggered for patients at high risk for PND (**Figure 2**). These delirium care pathways include a cautious approach to anesthesia that starts by avoiding deliriogenic medications and optimizing intraoperative anesthetic care. Postoperatively, these pathways also include healthy nutrition, delirium screening, and nonpharmacological protocols that emphasize patient orientation, early access to glasses and hearing aids, sleep-wake cycle regularity, promotion of mobility, and pain control (8, 58). While adopting these multidisciplinary interventions can be challenging, they can be feasibly performed in the perioperative setting (59, 60).

In summary, clinical perioperative brain health care has seen recent advancements in consistent nomenclature, consensus on evidence-based perioperative care pathways, and new tools that facilitate accurate and actionable assessment of cognition in clinical settings. More research is needed to elucidate the mechanisms of PNDs and how baseline vulnerabilities, including NCDs, affect postsurgical disease progression. Furthermore, gaining insights into new interventional targets to protect perioperative brain health is an active area of work and the focus of the remainder of this review.

VALIDITY OF ANIMAL MODELS FOR PNDs

Findings from animal models can be extended to the target clinical population if certain factors are satisfied (61). Construct validity, also referred to as etiologic validity, considers the relevance of methods that create the model to the factors that produce the clinical disease. For PNDs, the model should be triggered by aseptic trauma (mimicking the surgical intervention), usually within the context of heightened vulnerability, in which risk factors, which are similar to those that enhance the prevalence of the clinical condition, are introduced. Face validity considers whether the model recapitulates, with high fidelity, critical biochemical, neuropathological, and behavioral features of the disease. For the clinical features to be similar to those of PND, the assessment of animal behavior must reflect the clinical cognitive domains that are typically impaired; it is unlikely that a single animal behavioral assessment will suffice for the features that are evident in the heterogeneous syndrome of PNDs. Predictive or pharmacological validity is established when the model responds to treatments in a way that predicts the effects of those treatments in humans; once established, the model can be used as a tool for screening for efficacy of prospective interventions. We focus on models involving rats and mice as these are the most common species used for PND preclinical research (62).

PROVOKING THE PHENOTYPE IN ANIMAL MODELS OF PND

The surgical trauma must be accomplished aseptically, under a general anesthetic regimen that itself does not modify the pathogenesis, resulting in an injury that does not interfere with mobility that is required in a subsequent behavioral assessment of cognition. As sepsis itself can produce cognitive disturbances, possibly through molecular mechanisms that differ from those of aseptic trauma, it is crucial that surgery be performed under sterile conditions. Apart from asepsis, the surgical intervention must avoid other pathogenic pathways, such as ischemic-reperfusion injury, which itself can cause cognitive decline (63), possibly through a different mechanism. Some sedative hypnotics can affect cognitive function independent of the surgery; a notable class are the benzodiazepines, which can directly affect both neuroconnectivity (64) and immune responses (65) and confound the influence of trauma on these potential mechanisms of PND. Furthermore, as prolonged exposure to volatile anesthetic agents can itself produce cognitive disturbances (66), the surgical intervention needs to be of short duration when performed under general anesthesia with these agents. Because neither neuraxial anesthesia (67) nor the type of general anesthetic (68) has been shown to influence the risk for delirium or postoperative cognitive impairment, respectively, these features have not been explored in preclinical models. As it is unethical to produce surgical trauma without anesthesia, models that seek to determine the putative modulatory effects of anesthetics on postsurgical behavior require the use of surgical surrogates, such as damage-associated molecular patterns (DAMPs), to trigger the phenotype (69) and that can be administered to a nonanesthetized animal. Behavioral assays that measure attentiveness and/or memory usually require assessment in a freely moving animal; therefore, it is important to ascertain that the surgical intervention (especially one involving the musculoskeletal system) does not interfere with mobility.

ENHANCING THE VULNERABILITY IN ANIMAL MODELS OF PND

As advanced age is the dominant risk factor for the development of PNDs (2), older animals are frequently used in translational models (70) for construct validity; use of young adult rodents may still be useful to understand surgery-induced changes in normal animals as a comparator to define why risk factors such as age influence the development of this postoperative complication. In

the International Study of Postoperative Cognitive Dysfunction investigation, perioperative respiratory infection was identified as a significant risk factor (71); consequently, some investigators have incorporated respiratory infection with mycoplasma into their models (72). Surgical patients with metabolic syndrome have an increased risk for POD following either noncardiac (73) or cardiac surgery (74); similarly, posttraumatic cognitive decline was exacerbated in a rat model of metabolic syndrome (75). Some studies have found that obese surgical patients have more exaggerated PND (76); this observation has been replicated in animal models of diet-induced obesity (77). Perioperative disrupted sleep hygiene and/or obstructive sleep apnea increases the probability for developing POD (78), and this finding has been recapitulated in animal models of sleep deficiency (79). Postoperative administration of opiate narcotics increases the risk for POD (80) and the duration of memory impairments and can be included in the model for those reasons (81). There is increasing interest in the possibility that surgery may hasten the onset or progression of AD symptoms (82) and that similar postoperative neuropathological changes to those of AD are observed (83); these observations led to the use of transgenic animals to explore the pathological and behavioral changes that surgery induces in this AD-permissive genetic background in which the postoperative cognitive impairment is longer lasting than in wild-type animals (84). Modification of the intestinal microbiome may also increase vulnerability for posttraumatic cognitive impairment (85), and rectification of an unstable intestinal microbiome results in an improvement in cognitive function (86). Cancer has been shown to be associated with increased risk for the development of cognitive impairment in an animal model (87); this finding is now being studied clinically (88).

BEHAVIORAL ASSESSMENT OF COGNITIVE DOMAINS IN ANIMAL MODELS OF PND

The complex cognitive processes required to develop thoughts and produce actions in humans involve domains that include attention, perception, executive functions, learning, memory, and language; additionally, emotional and motivational processes influence cognitive performance (89). Rather than targeting the entire spectrum of cognition, investigators have sought assessments that quantify the impairments that are present in PNDs. POD is the most common PND; its clinical features include inattention, disorganized thinking, altered levels of consciousness, disorientation, memory impairment, perceptual disturbances, and psychomotor agitation or retardation. Of these clinical features of POD, only inattention and memory impairment can reliably be measured through behavioral assessment in rodents. The domain of executive function, incorporating the faculties of working memory and attention, is often dysfunctional in PNDs (90).

Attentiveness

The 5-choice serial reaction time task (5-CSRTT) is a robust method for measuring impairment in attentiveness that can be performed in both rats (91) and mice (92); this preclinical test was specifically developed to mirror the inattentiveness that is present in children with attention deficit hyperactivity disorder and has been shown to have both predictive (93) and construct validity (94). In this attentiveness paradigm, the rodent is required to perform a nose poke into one of five holes in response to a brief (2 s) flash of light from that hole; correct spatial location results in a food reward. Both accuracy and speed of stimulus discrimination define the attentional capacity to monitor a horizontal array of food-rewarding apertures. Training is performed over several days (typically involving a total of ~25 sessions), and animals that can successfully respond with 80% accuracy are subsequently used for the experiments. Indices that are monitored in the test situation include accuracy (percentage of correct responses), omission rate (failure to respond),

and latency to nose poke. Additionally, a distracting stimulus (e.g., noise) can be introduced to challenge attentional processes that are directed to the relevant visual stimulus. Limitations of the 5-CSRTT include a requirement for adequate visual sensory function and that the operant nature of the task in which early satiety (established by an increase in latency for the food pellet to be retrieved) may limit motivation.

Memory

Working memory in humans refers to the ability to represent information in mind and use it in service of our immediate goals. An example of a clinical test of this function is Digit Span, which requires the examinee to repeat back a string of digits in forward or backward order (31). Absent language as a parameter in animals, the tasks that are used involve memory in maze-like paradigms (95). The delayed Y-alternation test (96) exploits the tendency of rodents to choose an alternate, other than the originally explored, arm to enter on successive trials, with an intertrial delay of a specified time (usually seconds).

Episodic memory refers to recollection of what has been personally experienced, with a retention time that is longer than for working memory. In the Novel Object Recognition (NOR) test (97), the novelty-seeking curiosity of rodents is exploited. In an acquisitional session, the animal is confronted with two identical objects, and they devote approximately equal time to exploring each individual object. In the retention phase, the animal explores a situation in which one of the familiar objects is substituted with a novel object; the test requires the animal to discriminate between the novel and the familiar object. If the animal can recall the objects to which it was previously exposed, it spends significantly more time exploring the novel object; if episodic memory is impaired, the animal is unable to distinguish which is the novel object and spends approximately equal time with each. Care with respect to the nature of the objects, the discriminability of the stimulus, the length of the sessions, intertrial intervals, and environmental conditions must be exercised (98). The NOR test can also be reconfigured to measure working memory by shortening the retention interval (99).

The most common histopathological feature of preclinical models of PND is neuroinflammation, especially of the hippocampus, a critical region for spatial episodic memory in rodents and humans (100). Spatial memory behavioral tasks are usually performed within a maze; the Morris water maze (MWM), the Barnes maze (BM), and the radial arm maze (RAM) are discussed further because of their popularity in laboratories that use preclinical models of PND. In the MWM, visual cues surround a circular pool to guide the rodent to a submerged platform that is obscured by the opaque nature of the fluid when the animal is introduced into the pool from different starting locations (101). Because the wound from the surgical trauma must not be contaminated, trials in the pool cannot be undertaken before it has healed. Similarly, any limitation in swimming speed from the surgical trauma will confound the assessment. Similar spatial memory deficits were observed on the MWM in a transgenic mouse model of AD as those on a virtual MWM in patients with MCI due to AD, providing face validity evidence for this test (102). In the BM, the rodent is required to escape from a brightly lit circular platform by locating the correct hole (of the 20 that are radially arranged) that leads to a small dark recessed chamber (103). In the RAM, eight arms radiate from a circular central platform on which the animal is placed; the goal of the task is to select the arm that contains a food pellet (104). In the acquisition or learning phase for each of the spatial memory paradigms, the performance of the task is assessed by the time it takes to reach the platform, hole, or arm; in the retention phase of the task, the reference memory probe is assessed by the preference for a quadrant, hole, or arm when the platform, recessed chamber, or food pellet is removed.

For emotive (associative) memory, a form of implicit memory, a trace fear conditioning paradigm is used in which an unconditional stimulus (mild foot shock applied from the grid floor) is paired with either contextual or auditory cues in the acquisition phase (105). In the test phase of the task, placement of the rodent in the same context or subjected to the same auditory cue, but without the shock, provokes a fear response in which the animal freezes (no movement except for breathing) in anticipation of a foot shock. A decrease in the time spent freezing in response to an auditory cue indicates a lesion in the amygdala, while a lesion in the hippocampus limits freezing in response to the contextual cue.

DRUGGABLE TARGETS ESTABLISHED IN ANIMAL MODELS OF PNDs

Successful interventions for PNDs require a thorough understanding of the cellular and molecular species involved; studies in animal models have generated two plausible pathogenic mechanisms: inflammation and oxidative stress. In the following description of the pathogenesis of PND, the cellular and molecular targets for interventions with disease-modifying entities are identified. Symptom mitigation is not considered because of the difficulty of aligning postoperative clinical symptoms with animal models.

Inflammation

Based on the similarity of the innate immune response to trauma between surgical patients and animal models (106), as well as meta-analyses associating the degree of inflammation with outcomes in patients (107), both construct and face validities are satisfied. The dynamic nature of the inflammatory response to trauma requires consideration of temporal changes rather than exhaustively describing each immune-related cell or cytokine that is altered by surgery; without appropriately timing the intervention, its efficacy may not become evident, a problem that has plagued immunotherapy for dynamic diseases such as sepsis (108). Therefore, the stages of the immune response to surgical trauma are now considered (**Figure 3**).

Systemic inflammation is provoked by high molecular group box protein 1 (HMGB1), a DAMP that is passively released from physically disrupted traumatized tissue as it shuttles between the cytoplasmic and nuclear compartments, dictated by its posttranslational modifications. Subsequently, active, exocytotic release of cytosolic HMGB1 from immunocytes occurs under the influence of proinflammatory cytokines (109). Depending on its posttranslational redox state, released HMGB1 can bind up to 14 different pattern-recognition receptors, including the receptor for advanced glycation end products (RAGE) and Toll-like receptors (TLRs) on chemotactically attracted immunocytes [including circulating bone marrow-derived monocytes (BM-DMs)]. Through RAGE-dependent signaling pathways, the transcription factor nuclear factor κ B (NF- κ B) localizes to the nucleus, resulting in the upregulation of proinflammatory cytokine genes, including *TNF α* , *IL1 β* , and *IL6*; the cognate proteins of these genes peak in the circulation within hours of the trauma (110). Animal studies have revealed that neutralizing HMGB1 (69), tumor necrosis factor α (TNF α) (111), IL-1 β (110), or IL-6 (112) at this early stage can block the development of postoperative cognitive decline (**Table 1**); however, a thorough understanding of the effect of early immunotherapy on healing of traumatized tissue is required before considering these as therapeutic interventions. Glycyrrhizin, a natural triterpene glycoconjugate derived from the root of licorice, has been shown to bind to HMGB1 and to prevent proinflammatory cytokine activity (113) and postoperative cognitive decline (114); again, caution should be exercised before extending this finding to the clinical setting because of the possible adverse effects of glycyrrhizin on wound healing.




Figure 3

Inflammatory mechanisms involved in perioperative neurocognitive disorders. Aseptic trauma to the tibia provokes the release of high molecular group box protein 1 (HMGB1) from the traumatized tissues. HMGB1 binds to pattern-recognition receptors (PRRs) on circulating bone marrow–derived monocytes (BM-DMs), resulting in nuclear translocation of nuclear factor κ B (NF- κ B) and the subsequent upregulation of genes for proinflammatory cytokines and its release into the circulation; HMGB1 also upregulates the chemokine MCP-1. High levels of circulating proinflammatory cytokines can permeabilize the blood-brain barrier, promoting the influx of C-C chemokine receptor type 2 (CCR2)-expressing BM-DMs into the brain parenchyma via the chemoattraction of MCP-1 to CCR2 receptors. Under the influence of BM-DMs, resident quiescent microglia in the brain become activated; together, these tissue macrophages release proinflammatory cytokines and HMGB1, which can disrupt long-term potentiation (LTP), a synaptic plasticity mechanism required for learning and memory. Inflammation is resolved through efferents from the vagus nerve that inhibit NF- κ B nuclear translocation via the action of acetylcholine (ACh) and lipoxins/resolvins, which are mediated by the α 7 nicotinic acetylcholine receptor (α 7nAChR) and G protein–coupled receptors (GPCRs), respectively. Figure created by Dr. Caroline Prioleau and Sabrina Erlhoff.

The next stage is the transformation of systemic inflammation into neuroinflammation (Figure 3). Despite the obstacle presented by the blood-brain barrier (BBB) to unfettered passage of inflammatory molecules and immunocytes (115), the brain can still receive instruction from the innate immune response to peripheral injury through the circulating proinflammatory cytokines IL-1 β and TNF α , which can signal the central nervous system by binding to cognate receptors on either the afferent vagus nerve (116) or endothelial cells in the BBB (117) to provoke neuroinflammation. Additionally, the BBB can be permeabilized by high levels of circulating TNF α , which decreases cadherin and ZO-1, proteins that are involved in establishing the BBB's tight junctions (118). Another immunological mechanism for enhancing BBB permeability is through the activation of mast cells (119). Restoring the integrity of the BBB (and preventing postoperative neuroinflammation) can be accomplished with a mast cell stabilizer such as disodium cromoglycate (119) (Table 1). Other strategies designed to strengthen the BBB include blocking the angiotensin II receptor type 1 that is upregulated in the hippocampus postoperatively (120) and by neutralizing IL-17A (121), although such an intervention may adversely affect peripheral inflammation and wound healing (Table 1).

Through the permeabilized BBB, circulating BM-DMs translocate into the brain parenchyma after trauma (122), attracted by the chemokine MCP-1, which is upregulated by HMGB1 (69). The resident microglia sense the changes induced by cytokine signaling and the influx of BM-DMs and become activated (123), polarize to a proinflammatory state (M1), and release proinflammatory cytokines, which disrupt synaptic plasticity processes associated with learning and memory

Table 1 Table of druggable targets

Pathogenesis	Target	Intervention	Adverse event(s)	Reference
Inflammation	HMGB1	Monoclonal Ab	Poor healing	69
	IL-1 β	Receptor antagonist	Poor healing	110
	TNF α	Monoclonal Ab	Poor healing, infection	111
	IL-6	Tocilizumab	Poor healing	112
	HMGB1	Glycyrrhizin	Hypertension	114
	Mast cell, BBB	Sodium cromoglycate	Rash, joint pain, headache	119
	BBB, angiotensin II receptor 1	Candesartan	Cold and flu-like symptoms	120
	BBB, IL-17A	Secukinumab	Nasopharyngitis	121
	Bone marrow-derived monocytes	Clodrolip	Infection	122
	Microglia	PLX5622	Loss of CNS homeostasis	125
	Microglia	Mixed-lineage kinase 3 inhibitor	NK	127
	Microglia	Kv1.3 inhibitor	NK	128
	Microglia	Atorvastatin	Cold and flu-like symptoms	130
	Microglia	Pioglitazone	SOB, fatigue, micturition problems	131
	Increase BDNF	Cannabinoids	Psychotomimetic	133
	Increase BDNF	Exercise	NK	86
	A1 astrocytes	Cholecystokinin	Abdominal pain, flushing	136
	A1 astrocytes	Minocycline	GI problems, dizziness	137
	Resolution	AT-Rv1D1	NK	141
	Resolution	Maresin	NK	142
Resolution, oxidative stress	Omega-3 PUFAs	GI problems, back pain	143	
Resolution	α 7nAChR agonist	NK	146	
Resolution	Dexmedetomidine	Bradycardia, hypotension, sedation	149	
Oxidative Stress	COX-2	COX-2 inhibitor	Insomnia, GI problems, rhinitis	153

Abbreviations: α 7nAChR, α 7 nicotinic acetylcholine receptor; Ab, antibody; AT-Rv1D1, aspirin-triggered resolvin 1D1; BBB, blood-brain barrier; BDNF, brain-derived neurotrophic factor; COX, cyclooxygenase; CNS, central nervous system; GI, gastrointestinal; HMGB1, high molecular group box protein 1; IL, interleukin; NK, not known; PUFA, polyunsaturated fatty acid; SOB, shortness of breath; TNF α , tumor necrosis factor α .

(124); chemokines are also released, which attract more immunocytes, thereby prolonging neuroinflammation (125).

Notwithstanding the adverse consequences of neuroinflammation through interruption of synaptic plasticity, neurotransmission, and even neuroapoptosis, the cellular constituents may also be a mechanism for regeneration and repair, for example, through release of trophic substances such as brain-derived neurotrophic factor (BDNF) from the reparative M2 microglia phenotype (126). Thus, while elimination of microglia with an antagonist directed at the colony stimulating factor-1 receptor (PLX5622) can prevent postoperative cognitive decline (125), this is unlikely to be translatable into clinical utility because of the crucial surveillance function of microglia. Rather, strategies aimed at selectively preventing activation of microglia are more appropriate; for example, blocking the mixed-lineage kinase 3 (127) or the Kv1.3 ion channel (128) can prevent microglia activation and postoperative cognitive decline while leaving intact the peripheral inflammatory response and wound healing—an ideal combination (**Table 1**). Conversion of microglia into the M2

phenotype has been accomplished with peroxisome proliferator-activated receptor γ (PPAR γ) agonists in other brain injuries (129); pretreatment with the PPAR γ atorvastatin reduced activated microglia and cognitive impairment (130), while pretreatment with pioglitazone, another PPAR γ , prevented the development of the surgical phenotype and increased expression of BDNF (131) (**Table 1**). It is noteworthy that BDNF dysregulation features prominently in postoperative cognitive impairment (132), and lower levels are present in comorbidities (e.g., type 2 diabetes, obesity) that increase the risk for developing postoperative cognitive impairment. Acute administration of cannabinoids can increase levels of BDNF (133) and can prevent postoperative impairment through cannabinoid 2 receptor signaling (134). Interestingly, nonpharmacologic interventions that increase BDNF levels (e.g., exercise) also prevent postoperative cognitive impairment (86).

In much the same manner that activation of microglia can result in proinflammatory (M1) or proregenerative (M2) phenotypes, the A1 reactive astrocyte phenotype can lose its trophic function and become neurotoxic (135). In an aged mouse model of PND, surgery induces astrocytes to express complement 3, a marker of the A1 phenotype (136). Despite the suppressant effect of either minocycline (137) or cholecystokinin (136) on the A1 phenotype and postoperative cognitive impairment, the nonspecific nature of these interventions precludes assigning causality (**Table 1**).

The resolution phase of inflammation is an active process involving both humoral and neural mechanisms. Humoral factors include specialized proresolving mediators (SPMs) that are broadly classified as lipoxins, resolvins (D/E series), and protectins/maresins, which are biotransformation products of the fatty acids arachidonic acid, eicosapentaenoic acid (EPA), and docosahexaenoic acid (DHA) (138). As these fatty acid precursors can also give rise to proinflammatory mediators such as prostaglandins and leukotrienes, precise temporal coordination is required to toggle or class switch between proinflammatory and proresolving states depending on the phase of inflammation (139). In conditions such as diabetes and metabolic syndrome, and with advanced age, biotransformation processes do not appropriately class switch, resulting in a dysregulated inflammatory response that is both exaggerated and prolonged, as was noted postoperatively in a rat model of the metabolic syndrome (140). SPMs bind to G protein-coupled receptors on circulating lymphocytes and monocytes, as well as tissue macrophages, to limit inflammation through inhibition of NF- κ B, increased phagocytosis, efferocytosis, and conversion to reparative cellular phenotypes.

Pretreatments that increase levels of SPMs can attenuate inflammation and prevent postoperative cognitive impairment in PND models (**Table 1**). For example, administration of an aspirin-triggered D series resolvin (AT-RvD1) prevented cognitive decline *in vivo* and relieved the impediment to long-term potentiation in hippocampal brain slices *ex vivo* (141). Prophylactic administration of maresin decreased systemic inflammation, entry of BM-DMs into the hippocampus, and cognitive impairment while sustaining the proresolving cytokine IL-10 and wound healing (142). As both EPA and DHA are derived from omega-3 polyunsaturated fatty acids (PUFAs), administration of fish oil (containing omega-3 PUFAs) perioperatively (3 days before until 7 days after the operation) improved cognition (assessed by the MWM), reduced peripheral inflammation, and resisted oxidative stress injury (143).

Neural mechanisms for resolving inflammation are mediated by the release of acetylcholine—under the influence of postganglionic vagal efferents—which binds to $\alpha 7$ nicotinic acetylcholine receptors ($\alpha 7$ nAChRs) on circulating BM-DMs as they traverse the spleen (144); $\alpha 7$ nAChR signaling inhibits nuclear translocation of NF- κ B and proinflammatory cytokine production. Interruption of this pathway with anticholinergic drugs enhances inflammation and cognitive impairment (145). The cholinergic inflammation-resolving pathway has been exploited through direct vagal stimulation (146) or by administration of the synthetic $\alpha 7$ nAChR agonist (146, 147) to limit postoperative inflammation and cognitive impairment (**Table 1**). Dexmedetomidine,

a sedative hypnotic that limits postoperative cognitive decline in humans (148), produces its ameliorative action through its vagomimetic properties (149).

The neural and humoral inflammation-resolving processes may have in common the upregulation of netrin-1 (150), which transduces its resolving action through neogenin 1; thus far, no interventions have been shown to modulate the levels of these key proteins to facilitate resolution of inflammation.

Oxidative Stress

Elaboration of reactive oxygen species (ROS) and depletion of antioxidants (needed to neutralize the toxic intermediates) reflect oxidative stress, which occurs after tissue trauma and has been associated with cognitive decline (151). Peripheral oxidative stress can disrupt the BBB and thereby facilitate the development of neuroinflammation; additionally, central oxidative stress is induced by the release of ROS and nitric oxide (a reactive nitrogen species) from activated microglia. Therefore, it is unclear whether oxidative stress is a unique process, a consequence of the inflammatory response to trauma, or a feedforward mechanism for prolonging neuroinflammation (152). Cyclooxygenase (COX) inhibitors, particularly of the COX-2 isoform, can reduce postoperative cognitive decline in rats (153) and are used for the management of postoperative inflammatory pain, although their efficacy for ameliorating postoperative cognitive impairment in patients has yet to be reported (NCT02689024).

CONCLUSION

Recent advances in clinical and preclinical investigations of PND have led to improved patient care and new intervention targets. In the clinical field, achievements include the standardization of nomenclature in alignment with the DSM-5, consensus on evidence-based perioperative care pathways, and new tools that facilitate rapid and accurate assessment of cognition that can detect preoperative cognitive impairment for risk stratification as well as establish a baseline for subsequent changes in the cognitive trajectory. In validated animal models, promising cellular and molecular targets have been identified in the innate immune response to aseptic trauma that are ripe for testing interventions for both efficacy and safety. Via the translation of discoveries in the preclinical environment and more widespread implementation of brain health pathways, the recent developments outlined in this review provide hope for better protection of brain health for vulnerable adults undergoing surgery.

DISCLOSURE STATEMENT

O.L.B.C. served as a clinical investigator for the OLIVER clinical trial from Medtronic. K.L.P. received research funding from Quest Diagnostics. M.M. is a co-holder of the patent for the use of dexmedetomidine as a sedative agent; he has not and will not receive royalty payments for the sale of dexmedetomidine as a sedative agent.

ACKNOWLEDGMENTS

The authors wish to thank Dr. Caroline Prioleau and Sabrina Erlhoff for their expert help in designing each of the figures. O.L.B.C. received funding from the National Institute of Neurological Disorders and Stroke (3UH3NS10557-03S1) and the Weill Pilot Award for Junior Investigators in Neurosciences. K.L.P. received grants from the National Institute of Neurological Disorders and Stroke (3UH3NS10557), the National Institute on Aging, Global Brain Health Institute,

LITERATURE CITED

1. Fowler AJ, Abbott TEF, Prowle J, Pearse RM. 2019. Age of patients undergoing surgery. *Br. J. Surg.* 106:1012–18
2. Monk TG, Weldon BC, Garvan CW, Dede DE, van der Aa MT, et al. 2008. Predictors of cognitive dysfunction after major noncardiac surgery. *Anesthesiology* 108:18–30
3. Leslie DL, Marcantonio ER, Zhang Y, Leo-Summers L, Inouye SK. 2008. One-year health care costs associated with delirium in the elderly population. *Arch. Intern. Med.* 168:27–32
4. Gleason LJ, Schmitt EM, Kosar CM, Tabloski P, Saczynski JS, et al. 2015. Effect of delirium and other major complications on outcomes after elective surgery in older adults. *JAMA Surg.* 150:1134–40
5. Rudolph JL, Jones RN, Rasmussen LS, Silverstein JH, Inouye SK, Marcantonio ER. 2007. Independent vascular and cognitive risk factors for postoperative delirium. *Am. J. Med.* 120:807–13
6. Honda S, Furukawa K, Nishiwaki N, Fujiya K, Omori H, et al. 2018. Risk factors for postoperative delirium after gastrectomy in gastric cancer patients. *World J. Surg.* 42:3669–75
7. Am. Psychiatr. Assoc. 2013. Neurocognitive disorders. In *Diagnostic and Statistical Manual of Mental Disorders*. Washington, DC: Am. Psychiatr. Publ. 5th ed.
8. Berger M, Schenning KJ, Brown CH 4th, Deiner SG, Whittington RA, et al. 2018. Best practices for postoperative brain health: recommendations from the Fifth International Perioperative Neurotoxicity Working Group. *Anesth. Analg.* 127:1406–13
9. Evered L, Silbert B, Knopman DS, Scott DA, DeKosky ST, et al. 2018. Recommendations for the nomenclature of cognitive change associated with anaesthesia and surgery—2018. *Anesth. Analg.* 127:1189–95
10. Deiner S, Liu X, Lin HM, Jacoby R, Kim J, et al. 2021. Does postoperative cognitive decline result in new disability after surgery? *Ann. Surg.* 274:e1108–14
11. Rajan KB, Weuve J, Barnes LL, McAninch EA, Wilson RS, Evans DA. 2021. Population estimate of people with clinical Alzheimer's disease and mild cognitive impairment in the United States (2020–2060). *Alzheimers Dement.* 17:1966–75
12. Silbert B, Evered L, Scott DA, McMahan S, Choong P, et al. 2015. Preexisting cognitive impairment is associated with postoperative cognitive dysfunction after hip joint replacement surgery. *Anesthesiology* 122:1224–34
13. Greene NH, Attix DK, Weldon BC, Smith PJ, McDonagh DL, Monk TG. 2009. Measures of executive function and depression identify patients at risk for postoperative delirium. *Anesthesiology* 110:788–95
14. Lee HB, Mears SC, Rosenberg PB, Leoutsakos JM, Gottschalk A, Sieber FE. 2011. Predisposing factors for postoperative delirium after hip fracture repair in individuals with and without dementia. *J. Am. Geriatr. Soc.* 59:2306–13
15. Culley DJ, Flaherty D, Fahey MC, Rudolph JL, Javedan H, et al. 2017. Poor performance on a preoperative cognitive screening test predicts postoperative complications in older orthopedic surgical patients. *Anesthesiology* 127:765–74
16. Sachdev PS, Blacker D, Blazer DG, Ganguli M, Jeste DV, et al. 2014. Classifying neurocognitive disorders: the DSM-5 approach. *Nat. Rev. Neurol.* 10:634–42
17. Brenowitz WD, Hubbard RA, Keene CD, Hawes SE, Longstreth WT Jr., et al. 2017. Mixed neuropathologies and estimated rates of clinical progression in a large autopsy sample. *Alzheimers Dement.* 13:654–62
18. McKeith IG, Boeve BF, Dickson DW, Halliday G, Taylor JP, et al. 2017. Diagnosis and management of dementia with Lewy bodies: fourth consensus report of the DLB Consortium. *Neurology* 89:88–100
19. Am. Soc. Anesthesiol. 2022. Perioperative Brain Health Initiative. *American Society of Anesthesiologists*. <https://www.asahq.org/brainhealthinitiative>
20. Chow WB, Rosenthal RA, Merkow RP, Ko CY, Esnaola NF, et al. 2012. Optimal preoperative assessment of the geriatric surgical patient: a best practices guideline from the American College of Surgeons

- National Surgical Quality Improvement Program and the American Geriatrics Society. *J. Am. Coll. Surg.* 215:453–66
21. Mohanty S, Rosenthal RA, Russell MM, Neuman MD, Ko CY, Esnaola NF. 2016. Optimal perioperative management of the geriatric patient: a best practices guideline from the American College of Surgeons NSQIP and the American Geriatrics Society. *J. Am. Coll. Surg.* 222:930–47
 22. Connolly A, Gahl E, Martin H, Morris J, Purandare N. 2011. Underdiagnosis of dementia in primary care: variations in the observed prevalence and comparisons to the expected prevalence. *Ageing Ment. Health* 15:978–84
 23. Bradford A, Kunik ME, Schulz P, Williams SP, Singh H. 2009. Missed and delayed diagnosis of dementia in primary care: prevalence and contributing factors. *Alzheimer Dis. Assoc. Disord.* 23:306–14
 24. Kotagal V, Langa KM, Plassman BL, Fisher GG, Giordani BJ, et al. 2015. Factors associated with cognitive evaluations in the United States. *Neurology* 84:64–71
 25. Tsoy E, Kiekhoefer RE, Guterman EL, Tee BL, Windon CC, et al. 2021. Assessment of racial/ethnic disparities in timeliness and comprehensiveness of dementia diagnosis in California. *JAMA Neurol.* 78:657–65
 26. Evered L. 2019. Cognitive testing for perioperative neurocognitive disorder. In *The Perioperative Neurocognitive Disorders*, ed. RG Ekenhoff, N Terrando, pp. 123–33. Cambridge, UK: Cambridge Univ. Press
 27. Deiner S, Fleisher LA, Leung JM, Peden C, Miller T, et al. 2020. Adherence to recommended practices for perioperative anesthesia care for older adults among US anesthesiologists: results from the ASA Committee on Geriatric Anesthesia–Perioperative Brain Health Initiative ASA member survey. *Perioper. Med.* 9:6
 28. Patnode CD, Perdue LA, Rossom RC, Rushkin MC, Redmond N, et al. 2020. *Screening for Cognitive Impairment in Older Adults: An Evidence Update for the U.S. Preventive Services Task Force*. Rockville, MD: Agency Healthc. Res. Qual.
 29. Tsoy E, Sideman AB, Pina Escudero SD, Pintado-Caipa M, Kanjanapong S, et al. 2021. Global perspectives on brief cognitive assessments for dementia diagnosis. *J. Alzheimers Dis.* 82:1001–13
 30. Possin KL, Tsoy E, Windon CC. 2021. Perils of race-based norms in cognitive testing: the case of former NFL players. *JAMA Neurol.* 78:377–78
 31. Rabinovici GD, Stephens ML, Possin KL. 2015. Executive dysfunction. *Continuum* 21:646–59
 32. Matthews BR. 2015. Memory dysfunction. *Continuum* 21:613–26
 33. van Harten AC, Smits LL, Teunissen CE, Visser PJ, Koene T, et al. 2013. Preclinical AD predicts decline in memory and executive functions in subjective complaints. *Neurology* 81:1409–16
 34. Cavallari M, Dai W, Guttman CRG, Meier DS, Ngo LH, et al. 2017. Longitudinal diffusion changes following postoperative delirium in older people without dementia. *Neurology* 89:1020–27
 35. Tanabe S, Mohanty R, Lindroth H, Casey C, Ballweg T, et al. 2020. Cohort study into the neural correlates of postoperative delirium: the role of connectivity and slow-wave activity. *Br. J. Anaesth.* 125:55–66
 36. Possin KL, Moskowitz T, Erlhoff SJ, Rogers KM, Johnson ET, et al. 2018. The brain health assessment for detecting and diagnosing neurocognitive disorders. *J. Am. Geriatr. Soc.* 66:150–66
 37. Cavallari M, Dai W, Guttman CR, Meier DS, Ngo LH, et al. 2016. Neural substrates of vulnerability to postsurgical delirium as revealed by presurgical diffusion MRI. *Brain* 139:1282–94
 38. Huang C, Martensson J, Gogenur I, Asghar MS. 2018. Exploring postoperative cognitive dysfunction and delirium in noncardiac surgery using MRI: a systematic review. *Neural Plast.* 2018:1281657
 39. Tsoy E, Erlhoff SJ, Goode CA, Dorsman KA, Kanjanapong S, et al. 2020. BHA-CS: a novel cognitive composite for Alzheimer’s disease and related disorders. *Alzheimers Dement.* 12:e12042
 40. Alioto AG, Mumford P, Wolf A, Casaletto KB, Erlhoff S, et al. 2019. White matter correlates of cognitive performance on the UCSF Brain Health Assessment. *J. Int. Neuropsychol. Soc.* 25:654–58
 41. Tsoy E, Strom A, Iaccarino L, Erlhoff SJ, Goode CA, et al. 2021. Detecting Alzheimer’s disease biomarkers with a brief tablet-based cognitive battery: sensitivity to A β and tau PET. *Alzheimers Res. Ther.* 13:36
 42. Libon DJ, McMillan C, Gunawardena D, Powers C, Massimo L, et al. 2009. Neurocognitive contributions to verbal fluency deficits in frontotemporal lobar degeneration. *Neurology* 73:535–42

43. Rodriguez-Salgado AM, Llibre-Guerra JJ, Tsoy E, Penalver-Guia AI, Bringas G, et al. 2021. A brief digital cognitive assessment for detection of cognitive impairment in Cuban older adults. *J. Alzheimers Dis.* 79:85–94
44. Bernstein Sideman A, Chalmer R, Ayers E, Gershon R, Verghese J, et al. 2022. Lessons from Detecting Cognitive Impairment Including Dementia (DetectCID) in primary care. *J. Alzheimers Dis.* 86:655–65
45. Evered LA, Silbert BS. 2018. Postoperative cognitive dysfunction and noncardiac surgery. *Anesth. Analg.* 127:496–505
46. Evered L, Scott DA, Silbert B, Maruff P. 2011. Postoperative cognitive dysfunction is independent of type of surgery and anesthetic. *Anesth. Analg.* 112:1179–85
47. Newman MF, Kirchner JL, Phillips-Bute B, Gaver V, Grocott H, et al. 2001. Longitudinal assessment of neurocognitive function after coronary-artery bypass surgery. *N. Engl. J. Med.* 344:395–402
48. Staffaroni AM, Tsoy E, Taylor J, Boxer AL, Possin KL. 2020. Digital cognitive assessments for dementia: digital assessments may enhance the efficiency of evaluations in neurology and other clinics. *Pract. Neurol.* 2020:24–45
49. Petersen RC, Lopez O, Armstrong MJ, Getchius TSD, Ganguli M, et al. 2018. Practice guideline update summary: mild cognitive impairment: report of the Guideline Development, Dissemination, and Implementation Subcommittee of the American Academy of Neurology. *Neurology* 90:126–35
50. Neurology. 1994. Practice parameter for diagnosis and evaluation of dementia (summary statement): report of the Quality Standards Subcommittee of the American Academy of Neurology. *Neurology* 44:2203–6
51. Burke D, Sengoz A, Schwartz R. 2000. Potentially reversible cognitive impairment in patients presenting to a memory disorders clinic. *J. Clin. Neurosci.* 7:120–23
52. Atkins KJ, Scott DA, Silbert B, Pike KE, Evered L. 2021. Preventing delirium and promoting long-term brain health: a clinical trial design for the Perioperative Cognitive Enhancement (PROTECT) trial. *J. Alzheimers Dis.* 83:1637–49
53. Ishizawa Y. 2022. Does preoperative cognitive optimization improve postoperative outcomes in the elderly? *J. Clin. Med.* 11:445
54. Gillis C, Buhler K, Bresee L, Carli F, Gramlich L, et al. 2018. Effects of nutritional prehabilitation, with and without exercise, on outcomes of patients who undergo colorectal surgery: a systematic review and meta-analysis. *Gastroenterology* 155:391–410.e4
55. Whittle J, Wischmeyer PE, Grocott MPW, Miller TE. 2018. Surgical prehabilitation: nutrition and exercise. *Anesthesiol. Clin.* 36:567–80
56. Humeidan ML, Reyes JC, Mavarez-Martinez A, Roeth C, Nguyen CM, et al. 2021. Effect of cognitive prehabilitation on the incidence of postoperative delirium among older adults undergoing major noncardiac surgery: the Neurobics randomized clinical trial. *JAMA Surg.* 156:148–56
57. O’Gara BP, Mueller A, Gasangwa DVI, Patxot M, Shaefi S, et al. 2020. Prevention of early postoperative decline: a randomized, controlled feasibility trial of perioperative cognitive training. *Anesth. Analg.* 130:586–95
58. Hughes CG, Boncyk CS, Culley DJ, Fleisher LA, Leung JM, et al. 2020. American Society for Enhanced Recovery and Perioperative Quality initiative joint consensus statement on postoperative delirium prevention. *Anesth. Analg.* 130:1572–90
59. Decker J, Kaloostian CL, Gurvich T, Nguyen P, Widjaja W, et al. 2020. Beyond cognitive screening: establishing an interprofessional perioperative brain health initiative. *J. Am. Geriatr. Soc.* 68:2359–64
60. Inouye SK, Bogardus ST Jr., Baker DI, Leo-Summers L, Cooney LM Jr. 2000. The Hospital Elder Life Program: a model of care to prevent cognitive and functional decline in older hospitalized patients. *J. Am. Geriatr. Soc.* 48:1697–706
61. Belzung C, Lemoine M. 2011. Criteria of validity for animal models of psychiatric disorders: focus on anxiety disorders and depression. *Biol. Mood Anxiety Disord.* 1:9
62. Eckenhoff RG, Maze M, Xie Z, Culley DJ, Goodlin SJ, et al. 2020. Perioperative neurocognitive disorder: state of the preclinical science. *Anesthesiology* 132:55–68
63. Zhang L, Liu H, Jia L, Lyu J, Sun Y, et al. 2019. Exosomes mediate hippocampal and cortical neuronal injury induced by hepatic ischemia-reperfusion injury through activating pyroptosis in rats. *Oxid. Med. Cell. Longev.* 2019:3753485

64. Numan T, van Dellen E, Vleggaar FP, van Vlieberghe P, Stam CJ, Slooter AJC. 2019. Resting state EEG characteristics during sedation with midazolam or propofol in older subjects. *Clin. EEG Neurosci.* 50:436–43
65. Sanders RD, Grover V, Goulding J, Godlee A, Gurney S, et al. 2015. Immune cell expression of GABA_A receptors and the effects of diazepam on influenza infection. *J. Neuroimmunol.* 282:97–103
66. Culley DJ, Baxter MG, Crosby CA, Yukhananov R, Crosby G. 2004. Impaired acquisition of spatial memory 2 weeks after isoflurane and isoflurane-nitrous oxide anesthesia in aged rats. *Anesth. Analg.* 99:1393–97
67. Neuman MD, Feng R, Carson JL, Gaskins LJ, Dillane D, et al. 2021. Spinal anesthesia or general anesthesia for hip surgery in older adults. *N. Engl. J. Med.* 385:2025–35
68. Jiao XF, Lin XM, Ni XF, Li HL, Zhang C, et al. 2019. Volatile anesthetics versus total intravenous anesthesia in patients undergoing coronary artery bypass grafting: an updated meta-analysis and trial sequential analysis of randomized controlled trials. *PLOS ONE* 14:e0224562
69. Vacas S, Degos V, Tracey KJ, Maze M. 2014. High-mobility group box 1 protein initiates postoperative cognitive decline by engaging bone marrow-derived macrophages. *Anesthesiology* 120:1160–67
70. Li Z, Liu F, Ma H, White PF, Yumul R, et al. 2017. Age exacerbates surgery-induced cognitive impairment and neuroinflammation in Sprague-Dawley rats: the role of IL-4. *Brain Res.* 1665:65–73
71. Moller JT, Cluitmans P, Rasmussen LS, Houx P, Rasmussen H, et al. 1998. Long-term postoperative cognitive dysfunction in the elderly: ISPOCD1 study. *Lancet* 351:857–61
72. Hovens IB, van Leeuwen BL, Nyakas C, Heineman E, van der Zee EA, Schoemaker RG. 2015. Prior infection exacerbates postoperative cognitive dysfunction in aged rats. *Am. J. Physiol. Regul. Integr. Comp. Physiol.* 309:R148–59
73. Hudetz JA, Patterson KM, Amole O, Riley AV, Pagel PS. 2011. Postoperative cognitive dysfunction after noncardiac surgery: effects of metabolic syndrome. *J. Anesth.* 25:337–44
74. Hudetz JA, Patterson KM, Iqbal Z, Gandhi SD, Pagel PS. 2011. Metabolic syndrome exacerbates short-term postoperative cognitive dysfunction in patients undergoing cardiac surgery: results of a pilot study. *J. Cardiothorac. Vasc. Anesth.* 25:282–87
75. Feng X, Degos V, Koch LG, Britton SL, Zhu Y, et al. 2013. Surgery results in exaggerated and persistent cognitive decline in a rat model of the metabolic syndrome. *Anesthesiology* 118:1098–105
76. Feinkohl I, Winterer G, Pischon T. 2016. Obesity and post-operative cognitive dysfunction: a systematic review and meta-analysis. *Diabetes Metab. Res. Rev.* 32:643–51
77. Wei L, Yao M, Zhao Z, Jiang H, Ge S. 2018. High-fat diet aggravates postoperative cognitive dysfunction in aged mice. *BMC Anesthesiol.* 18:20
78. Todd OM, Gelrich L, MacLulich AM, Driessen M, Thomas C, Kreisel SH. 2017. Sleep disruption at home as an independent risk factor for postoperative delirium. *J. Am. Geriatr. Soc.* 65:949–57
79. Ni P, Dong H, Zhou Q, Wang Y, Sun M, et al. 2019. Preoperative sleep disturbance exaggerates surgery-induced neuroinflammation and neuronal damage in aged mice. *Mediators Inflamm.* 2019:8301725
80. Shehabi Y, Grant P, Wolfenden H, Hammond N, Bass F, et al. 2009. Prevalence of delirium with dexmedetomidine compared with morphine based therapy after cardiac surgery: a randomized controlled trial (DEXmedetomidine COMpared to Morphine—DEXCOM study). *Anesthesiology* 111:1075–84
81. Muscat SM, Deems NP, D'Angelo H, Kitt MM, Grace PM, et al. 2021. Postoperative cognitive dysfunction is made persistent with morphine treatment in aged rats. *Neurobiol. Aging* 98:214–24
82. Silverstein JH. 2014. Influence of anesthetics on Alzheimer's disease: biophysical, animal model, and clinical reports. *J. Alzheimers Dis.* 40:839–48
83. Shen W, Lu K, Wang J, Wu A, Yue Y. 2016. Activation of mTOR signaling leads to orthopedic surgery-induced cognitive decline in mice through β -amyloid accumulation and tau phosphorylation. *Mol. Med. Rep.* 14:3925–34
84. Tang JX, Mardini F, Janik LS, Garrity ST, Li RQ, et al. 2013. Modulation of murine Alzheimer pathogenesis and behavior by surgery. *Ann. Surg.* 257:439–48
85. Zhan G, Hua D, Huang N, Wang Y, Li S, et al. 2019. Anesthesia and surgery induce cognitive dysfunction in elderly male mice: the role of gut microbiota. *Aging* 11:1778–90

86. Feng X, Uchida Y, Koch L, Britton S, Hu J, et al. 2017. Exercise prevents enhanced postoperative neuroinflammation and cognitive decline and rectifies the gut microbiome in a rat model of metabolic syndrome. *Front. Immunol.* 8:1768
87. Wu T, Wang X, Zhang R, Jiao Y, Yu W, et al. 2020. Mice with pre-existing tumors are vulnerable to postoperative cognitive dysfunction. *Brain Res.* 1732:146650
88. Zhang Y, Li HJ, Wang DX, Jia HQ, Sun XD, et al. 2017. Impact of inhalational versus intravenous anaesthesia on early delirium and long-term survival in elderly patients after cancer surgery: study protocol of a multicentre, open-label, and randomised controlled trial. *BMJ Open* 7:e018607
89. Al Dahhan NZ, De Felice FG, Munoz DP. 2019. Potentials and pitfalls of cross-translational models of cognitive impairment. *Front. Behav. Neurosci.* 13:48
90. Valentin LS, Pereira VF, Pietrobon RS, Schmidt AP, Oses JP, et al. 2016. Effects of single low dose of dexamethasone before noncardiac and nonneurologic surgery and general anesthesia on postoperative cognitive dysfunction—a Phase III double blind, randomized clinical trial. *PLOS ONE* 11:e0152308
91. Bari A, Dalley JW, Robbins TW. 2008. The application of the 5-choice serial reaction time task for the assessment of visual attentional processes and impulse control in rats. *Nat. Protoc.* 3:759–67
92. Patel S, Stolerman IP, Asherson P, Sluyter F. 2006. Attentional performance of C57BL/6 and DBA/2 mice in the 5-choice serial reaction time task. *Behav. Brain Res.* 170:197–203
93. Paine TA, Tomasiewicz HC, Zhang K, Carlezon WA Jr. 2007. Sensitivity of the five-choice serial reaction time task to the effects of various psychotropic drugs in Sprague-Dawley rats. *Biol. Psychiatry* 62:687–93
94. Velagapudi R, Subramaniam S, Xiong C, Porkka F, Rodriguiz RM, et al. 2019. Orthopedic surgery triggers attention deficits in a delirium-like mouse model. *Front. Immunol.* 10:2675
95. Dudchenko PA. 2004. An overview of the tasks used to test working memory in rodents. *Neurosci. Biobehav. Rev.* 28:699–709
96. Yang ST, Shi Y, Wang Q, Peng JY, Li BM. 2014. Neuronal representation of working memory in the medial prefrontal cortex of rats. *Mol. Brain* 7:61
97. Ennaceur A, Delacour J. 1988. A new one-trial test for neurobiological studies of memory in rats. 1: Behavioral data. *Behav. Brain Res.* 31:47–59
98. Darcet F, Gardier AM, Gaillard R, David DJ, Guilloux JP. 2016. Cognitive dysfunction in major depressive disorder. A translational review in animal models of the disease. *Pharmaceuticals* 9:9
99. Antunes M, Biala G. 2012. The novel object recognition memory: neurobiology, test procedure, and its modifications. *Cogn. Process.* 13:93–110
100. Bannerman DM, Sprengel R, Sanderson DJ, McHugh SB, Rawlins JN, et al. 2014. Hippocampal synaptic plasticity, spatial memory and anxiety. *Nat. Rev. Neurosci.* 15:181–92
101. Tucker LB, Velosky AG, McCabe JT. 2018. Applications of the Morris water maze in translational traumatic brain injury research. *Neurosci. Biobehav. Rev.* 88:187–200
102. Possin KL, Sanchez PE, Anderson-Bergman C, Fernandez R, Kerchner GA, et al. 2016. Cross-species translation of the Morris maze for Alzheimer's disease. *J. Clin. Investig.* 126:779–83
103. Pitts MW. 2018. Barnes maze procedure for spatial learning and memory in mice. *Bio Protoc.* 8:e2744
104. Mei J, Kohler J, Winter Y, Spies C, Endres M, et al. 2020. Automated radial 8-arm maze: a voluntary and stress-free behavior test to assess spatial learning and memory in mice. *Behav. Brain Res.* 381:112352
105. Curzon P, Rustay NR, Browman KE. 2009. Cued and contextual fear conditioning for rodents. In *Methods of Behavior Analysis in Neuroscience*, ed. JJ Buccafusco. Boca Raton, FL: CRC Press. 2nd ed.
106. Berger M, Murdoch DM, Staats JS, Chan C, Thomas JP, et al. 2019. Flow cytometry characterization of cerebrospinal fluid monocytes in patients with postoperative cognitive dysfunction: a pilot study. *Anesth. Analg.* 129:e150–54
107. Noah AM, Almghairbi D, Evley R, Moppett IK. 2021. Preoperative inflammatory mediators and postoperative delirium: systematic review and meta-analysis. *Br. J. Anaesth.* 127:424–34
108. Beckmann N, Salyer CE, Crisologo PA, Nomellini V, Caldwell CC. 2020. Staging and personalized intervention for infection and sepsis. *Surg. Infect.* 21:732–44
109. Saxena S, Krays V, De Jongh R, Vamecq J, Maze M. 2021. High-mobility group box-1 and its potential role in perioperative neurocognitive disorders. *Cells* 10:2582
110. Cibelli M, Fidalgo AR, Terrando N, Ma D, Monaco C, et al. 2010. Role of interleukin-1 β in postoperative cognitive dysfunction. *Ann. Neurol.* 68:360–68

111. Terrando N, Monaco C, Ma D, Foxwell BM, Feldmann M, Maze M. 2010. Tumor necrosis factor- α triggers a cytokine cascade yielding postoperative cognitive decline. *PNAS* 107:20518–22
112. Hu J, Feng X, Valdearcos M, Lutrin D, Uchida Y, et al. 2018. Interleukin-6 is both necessary and sufficient to produce perioperative neurocognitive disorder in mice. *Br. J. Anaesth.* 120:537–45
113. Mollica L, De Marchis F, Spitaleri A, Dallacosta C, Pennacchini D, et al. 2007. Glycyrrhizin binds to high-mobility group box 1 protein and inhibits its cytokine activities. *Chem. Biol.* 14:431–41
114. Kong ZH, Chen X, Hua HP, Liang L, Liu LJ. 2017. The oral pretreatment of glycyrrhizin prevents surgery-induced cognitive impairment in aged mice by reducing neuroinflammation and Alzheimer's-related pathology via HMGB1 inhibition. *J. Mol. Neurosci.* 63:385–95
115. Obermeier B, Daneman R, Ransohoff RM. 2013. Development, maintenance and disruption of the blood-brain barrier. *Nat. Med.* 19:1584–96
116. Steinberg BE, Silverman HA, Robbiati S, Gunasekaran MK, Tsaava T, et al. 2016. Cytokine-specific neurograms in the sensory vagus nerve. *Bioelectron. Med.* 3:7–17
117. Salvador AF, de Lima KA, Kipnis J. 2021. Neuromodulation by the immune system: a focus on cytokines. *Nat. Rev. Immunol.* 21:526–41
118. Lopez-Ramirez MA, Fischer R, Torres-Badillo CC, Davies HA, Logan K, et al. 2012. Role of caspases in cytokine-induced barrier breakdown in human brain endothelial cells. *J. Immunol.* 189:3130–39
119. Zhang S, Dong H, Zhang X, Li N, Sun J, Qian Y. 2016. Cerebral mast cells contribute to postoperative cognitive dysfunction by promoting blood brain barrier disruption. *Behav. Brain Res.* 298:158–66
120. Li Z, Mo N, Li L, Cao Y, Wang W, et al. 2016. Surgery-induced hippocampal angiotensin II elevation causes blood-brain barrier disruption via MMP/TIMP in aged rats. *Front. Cell Neurosci.* 10:105
121. Ni P, Dong H, Wang Y, Zhou Q, Xu M, et al. 2018. IL-17A contributes to perioperative neurocognitive disorders through blood-brain barrier disruption in aged mice. *J. Neuroinflamm.* 15:332
122. Degos V, Vacas S, Han Z, van Rooijen N, Gressens P, et al. 2013. Depletion of bone marrow-derived macrophages perturbs the innate immune response to surgery and reduces postoperative memory dysfunction. *Anesthesiology* 118:527–36
123. Hovens IB, van Leeuwen BL, Nyakas C, Heineman E, van der Zee EA, Schoemaker RG. 2015. Postoperative cognitive dysfunction and microglial activation in associated brain regions in old rats. *Neurobiol. Learn. Mem.* 118:74–79
124. Kelly A, Vereker E, Nolan Y, Brady M, Barry C, et al. 2003. Activation of p38 plays a pivotal role in the inhibitory effect of lipopolysaccharide and interleukin-1 β on long term potentiation in rat dentate gyrus. *J. Biol. Chem.* 278:19453–62
125. Feng X, Valdearcos M, Uchida Y, Lutrin D, Maze M, Koliwad SK. 2017. Microglia mediate postoperative hippocampal inflammation and cognitive decline in mice. *JCI Insight* 2:e91229
126. Zou J, Huang GF, Xia Q, Li X, Shi J, Sun N. 2022. Electroacupuncture promotes microglial M2 polarization in ischemic stroke via annexin A1. *Acupunct. Med.* 40:258–67
127. Miller-Rhodes P, Kong C, Baht GS, Saminathan P, Rodriguiz RM, et al. 2019. The broad spectrum mixed-lineage kinase 3 inhibitor URMC-099 prevents acute microgliosis and cognitive decline in a mouse model of perioperative neurocognitive disorders. *J. Neuroinflamm.* 16:193
128. Lai IK, Valdearcos M, Morioka K, Saxena S, Feng X, et al. 2020. Blocking Kv1.3 potassium channels prevents postoperative neuroinflammation and cognitive decline without impairing wound healing in mice. *Br. J. Anaesth.* 125:298–307
129. Lee CH, Park OK, Yoo KY, Byun K, Lee B, et al. 2011. The role of peroxisome proliferator-activated receptor γ , and effects of its agonist, rosiglitazone, on transient cerebral ischemic damage. *J. Neurol. Sci.* 300:120–29
130. Vizcaychipi MP, Watts HR, O'Dea KP, Lloyd DG, Penn JW, et al. 2014. The therapeutic potential of atorvastatin in a mouse model of postoperative cognitive decline. *Ann. Surg.* 259:1235–44
131. Zhang Z, Yuan H, Zhao H, Qi B, Li F, An L. 2017. PPAR γ activation ameliorates postoperative cognitive decline probably through suppressing hippocampal neuroinflammation in aged mice. *Int. Immunopharmacol.* 43:53–61
132. Qiu LL, Pan W, Luo D, Zhang GF, Zhou ZQ, et al. 2020. Dysregulation of BDNF/TrkB signaling mediated by NMDAR/Ca²⁺/calpain might contribute to postoperative cognitive dysfunction in aging mice. *J. Neuroinflamm.* 17:23

133. Ferreira FF, Ribeiro FF, Rodrigues RS, Sebastiao AM, Xapelli S. 2018. Brain-derived neurotrophic factor (BDNF) role in cannabinoid-mediated neurogenesis. *Front. Cell Neurosci.* 12:441
134. Sun L, Dong R, Xu X, Yang X, Peng M. 2017. Activation of cannabinoid receptor type 2 attenuates surgery-induced cognitive impairment in mice through anti-inflammatory activity. *J. Neuroinflamm.* 14:138
135. Liddelow SA, Guttenplan KA, Clarke LE, Bennett FC, Bohlen CJ, et al. 2017. Neurotoxic reactive astrocytes are induced by activated microglia. *Nature* 541:481–87
136. Chen L, Yang N, Li Y, Li Y, Hong J, et al. 2021. Cholecystokinin octapeptide improves hippocampal glutamatergic synaptogenesis and postoperative cognition by inhibiting induction of A1 reactive astrocytes in aged mice. *CNS Neurosci. Ther.* 27:1374–84
137. Jin WJ, Feng SW, Feng Z, Lu SM, Qi T, Qian YN. 2014. Minocycline improves postoperative cognitive impairment in aged mice by inhibiting astrocytic activation. *NeuroReport* 25:1–6
138. Chiang N, Serhan CN. 2020. Specialized pro-resolving mediator network: an update on production and actions. *Essays Biochem.* 64:443–62
139. Serhan CN. 2014. Pro-resolving lipid mediators are leads for resolution physiology. *Nature* 510:92–101
140. Su X, Feng X, Terrando N, Yan Y, Chawla A, et al. 2013. Dysfunction of inflammation-resolving pathways is associated with exaggerated postoperative cognitive decline in a rat model of the metabolic syndrome. *Mol. Med.* 18:1481–90
141. Terrando N, Gomez-Galan M, Yang T, Carlstrom M, Gustavsson D, et al. 2013. Aspirin-triggered resolvin D1 prevents surgery-induced cognitive decline. *FASEB J.* 27:3564–71
142. Yang T, Xu G, Newton PT, Chagin AS, Mkrtchian S, et al. 2019. Maresin 1 attenuates neuroinflammation in a mouse model of perioperative neurocognitive disorders. *Br. J. Anaesth.* 122:350–60
143. Guo Y, Ping F, Cao Y, Shang J, Zhang J, et al. 2021. Effects of omega-3 polyunsaturated fatty acids on cognitive function after splenectomy in rats. *Biomed. Res. Int.* 2021:5513886
144. Gallowitsch-Puerta M, Tracey KJ. 2005. Immunologic role of the cholinergic anti-inflammatory pathway and the nicotinic acetylcholine $\alpha 7$ receptor. *Ann. N. Y. Acad. Sci.* 1062:209–19
145. Terrando N, Eriksson LI, Ryu JK, Yang T, Monaco C, et al. 2011. Resolving postoperative neuroinflammation and cognitive decline. *Ann. Neurol.* 70:986–95
146. Xiong J, Wang H, Bao Y, Guo Y, Sun Y. 2019. Electric vagal nerve stimulation inhibits inflammation and improves early postoperation cognitive dysfunction in aged rats. *BMC Anesthesiol.* 19:217
147. Terrando N, Yang T, Ryu JK, Newton PT, Monaco C, et al. 2015. Stimulation of the $\alpha 7$ nicotinic acetylcholine receptor protects against neuroinflammation after tibia fracture and endotoxemia in mice. *Mol. Med.* 20:667–75
148. Su X, Meng ZT, Wu XH, Cui F, Li HL, et al. 2016. Dexmedetomidine for prevention of delirium in elderly patients after non-cardiac surgery: a randomised, double-blind, placebo-controlled trial. *Lancet* 388:1893–902
149. Hu J, Vacas S, Feng X, Lutrin D, Uchida Y, et al. 2018. Dexmedetomidine prevents cognitive decline by enhancing resolution of high mobility group box 1 protein-induced inflammation through a vagomimetic action in mice. *Anesthesiology* 128:921–31
150. Mirakaj V, Dalli J, Granja T, Rosenberger P, Serhan CN. 2014. Vagus nerve controls resolution and pro-resolving mediators of inflammation. *J. Exp. Med.* 211:1037–48
151. Robillard JM, Gordon GR, Choi HB, Christie BR, MacVicar BA. 2011. Glutathione restores the mechanism of synaptic plasticity in aged mice to that of the adult. *PLoS ONE* 6:e20676
152. Rojo AI, McBean G, Cindric M, Egea J, Lopez MG, et al. 2014. Redox control of microglial function: molecular mechanisms and functional significance. *Antioxid. Redox. Signal.* 21:1766–801
153. Wang YB, Chen Z, Li J, Shi J. 2019. Parecoxib improves the cognitive function of POCD rats via attenuating COX-2. *Eur. Rev. Med. Pharmacol. Sci.* 23:4971–79